



ORAL PRESENTATION

Open Access

No lack of regulatory B cells in patients with Multiple Sclerosis

Laure Michel^{1,2,3*}, Mélanie Chesneau^{1,3}, Philippe Manceau¹, Alexandra Garcia^{1,3}, Marion Salou^{1,3}, Annie Elong Ngonu^{1,3}, Annaïck Pallier^{1,3}, Marylène Jacq-Foucher², Fabienne Lefrère², Sandrine Wiertlewski², Jean-Paul Soullillou^{1,3}, Nicolas Degauque^{1,3}, David-Axel Laplaud^{1,2,3†}, Sophie Brouard^{1,3†}

From 7th European Workshop on Immune-Mediated Inflammatory Diseases Noordwijk aan Zee, the Netherlands. 28-30 November 2012

Background

Recent data support a prominent role for B cells in MS physiopathology. Recently it has emerged that subsets of B cells secreting IL-10 negatively regulate disease symptoms in Experimental Autoimmune Encephalomyelitis (EAE). However, the involvement of such regulatory B cells in MS remains unclear.

Aim

We aimed to study the frequency, phenotype and function of regulatory B cells in MS patients as compared to Healthy Volunteers (HV).

Methods

Sixty-three untreated MS patients and 58 HV were included in this study. IL-10 secretion by B cells and phenotype of IL-10⁺ B cells were studied after 5h (B10 cells) and 48h of stimulation (B10pro cells) by CD40L and ODN. Coculture assays with prestimulated B cells and responding CD4⁺CD25⁻ T cells were performed for 3 days.

Results

No significant difference was found either for IL-10 secretion ability of B cells after 5h or 48h of stimulation. The analysis of B10pro cells phenotype revealed mainly a memory phenotype in MS and HV, even if both naïve and immature subsets were also able to secrete IL-10. Prestimulated B cells from MS inhibited CD4⁺CD25⁻ T cell proliferation in the same manner than HV by a contact dependent mechanism, independently of IL-10 and TGF- β secretion.

Conclusion

Altogether, our data show that regulatory B cells have a conserved frequency, phenotype and function in the blood of patients with MS suggesting that B cells do not contribute to the physiopathology of the disease.

Acknowledgements

This work was funded by ARSEP.

Author details

¹INSERM, UMR 1064, Nantes, France. ²CHU Nord Laennec, Service de Neurologie, Nantes, France. ³CHU de Nantes, ITUN, Nantes, France.

Published: 28 November 2012

doi:10.1186/1479-5876-10-S3-O4

Cite this article as: Michel et al.: No lack of regulatory B cells in patients with Multiple Sclerosis. *Journal of Translational Medicine* 2012 **10**(Suppl 3):O4.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



† Contributed equally

¹INSERM, UMR 1064, Nantes, France

Full list of author information is available at the end of the article