



POSTER PRESENTATION

Open Access

Apolipoprotein E and undercarboxylated osteocalcin are associated with bone fragility but not with bone mineral density in osteoarthritis patients

Ana Rodrigues^{1,2*}, Joana Caetano-Lopes¹, Ana Lopes¹, Ana C Vale³, Inês Aleixo¹, Inês P Perpétuo¹, Ana S Pena¹, Alexandra Faustino¹, Alexandre Sepriano¹, Joaquim Polido-Pereira^{1,2}, Elsa Vieira-Sousa^{1,2}, Bruno Vidal¹, José C Romeu², Pedro M Amaral³, Luís G Rosa³, José A Pereira da Silva², Jacinto Monteiro⁴, Maria F Vaz³, João E Fonseca^{1,2}, Helena Canhão^{1,2}

From 6th European Workshop on Immune-Mediated Inflammatory Diseases
Nice, France. 23-25 November 2011

Background

Apolipoprotein E (apoE) gene polymorphisms and undercarboxylated osteocalcin (ucOC) and vitamin K have been associated with fragility fractures and low BMD in general population. The aim of this work was to study whether the effect of apoE gene polymorphisms, seric apoE levels and ucOC influence trabecular bone biomechanics and bone mineral density (BMD) in patients submitted to hip replacement surgery due to advanced primary osteoarthritis.

Materials and methods

Patients were evaluated for established clinical risk factors (CRFs) for fracture. A dual X-ray absorptiometry (DXA) was performed. ApoE genotyping was performed at positions rs429358 and rs7412. Fasting blood samples were collected at the time of surgery to assess the following parameters: seric apoE, ucOC, Vitamin K, LDL cholesterol, triglycerides and bone turnover markers.

Femoral epiphyses were collected and trabecular bone cylinders were drilled in order to perform compression mechanical tests and analyze bone strength and stiffness.

Results

Forty-four patients were studied (median 70 years of age, 55% women and 90% postmenopausal, BMI of 27.2

Kg/m²). 6.8% had prevalent fragility fractures, 36% had normal BMD and 64% were osteopenic. The apoE genotype distribution was in accordance with Hardy-Weinberg equilibrium and the E4 allele, previously documented as the risk allele, was present in 16% of the patients. This allele was significantly associated with lower trabecular strength ($p=0.004$) and stiffness ($p=0.07$), adjusted for age and gender, but not with BMD. Moreover, E4 allele was associated with higher levels of markers of bone formation: ALP ($p=0.067$), BSALP ($p=0.085$) and P1NP ($p=0.062$). We also found that ucOC was negatively correlated with strength ($p=0.077$) and stiffness ($p=0.047$), regardless of patients gender, age or vitamin K, but not with BMD. ApoE levels were also associated with strength ($p=0.056$; $R=-0.322$) and stiffness ($p=0.001$; $R=-0.513$).

Conclusions

Other studies have shown that advanced hip osteoarthritis can be associated with a higher BMD, but is not a protective factor for fragility fractures. Our observations suggest that in osteoarthritis patients Apo E4 allele, seric apo E and ucOC are biologic risk factors for fragility in a more independent way from BMD than in the general population.

Author details

¹Rheumatology Research Unit, Instituto de Medicina Molecular, Faculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal. ²Rheumatology and Bone Metabolic Diseases Dept., Hospital de Santa Maria, Lisbon, Portugal.

¹Rheumatology Research Unit, Instituto de Medicina Molecular, Faculdade de Medicina da Universidade de Lisboa, Lisbon, Portugal
Full list of author information is available at the end of the article

³Dept. of Mechanical Engineering, Instituto Superior Técnico, Lisbon, Portugal. ⁴Ortopaedics Dept., Hospital de Santa Maria, Lisbon, Portugal.

Published: 23 November 2011

doi:10.1186/1479-5876-9-S2-P54

Cite this article as: Rodrigues *et al.*: Apolipoprotein E and undercarboxylated osteocalcin are associated with bone fragility but not with bone mineral density in osteoarthritis patients. *Journal of Translational Medicine* 2011 **9**(Suppl 2):P54.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

