CORRECTION

Open Access



Correction to: Accumulation of blood-circulating PD-L1-expressing M-MDSCs and monocytes/macrophages in pretreatment ovarian cancer patients is associated with soluble PD-L1

Karolina Okła^{1*}, Alicja Rajtak¹, Arkadiusz Czerwonka², Marcin Bobiński¹, Anna Wawruszak³, Rafał Tarkowski¹, Wiesława Bednarek¹, Justyna Szumiło⁴ and Jan Kotarski¹

Correction to: J Transl Med (2020) 18:220

https://doi.org/10.1186/s12967-020-02389-7 Following the publication of the original article [1], it was noted that due to a typesetting error, the Fig. 1 was replaced by a duplicate of Fig. 3. The correct Fig. 1 is given below, and the original article has been corrected.


The original article can be found online at https://doi.org/10.1186/s1296 7-020-02389-7

*Correspondence: karolinaokla@umlub.pl ¹ The First Department of Oncologic Gynecology and Gynecology, Medical University of Lublin, 20-081 Lublin, Poland Full list of author information is available at the end of the article



© The Author(s) 2020. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.





(MO/MA). Simultaneous analysis of myeloid cell populations in the **a**, **b** blood, **c** ascites and **d** tumour tissue of patients with ovarian cancer. Mononuclear cells (MCs) obtained from the blood (n = 43), ascites (n = 26), and tumour tissue (n = 29) of ovarian cancer patients and from the blood of healthy women (n = 15) were analyzed using flow cytometry. The levels of M-MDSCs and MO/MA are presented as the percentage of MCs. The horizontal lines are the median values and the whiskers indicate the minimum and maximum values. Each point corresponds to an individual patient. *p < 0.05; ***p < 0.001; ****p < 0.001

Author details

¹ The First Department of Oncologic Gynecology and Gynecology, Medical University of Lublin, 20-081 Lublin, Poland. ² Department of Virology and Immunology, Maria Curie-Sklodowska University, 20-031, Lublin, Poland. ³ Department of Biochemistry and Molecular Biology, Medical University of Lublin, 20-081 Lublin, Poland. ⁴ Department of Clinical Pathomorphology, Medical University of Lublin, 20-090 Lublin, Poland.

Published online: 25 June 2020

Reference

 Okła K, Rajtak A, Czerwonka A, Bobiński M, Wawruszak A, Tarkowski R, Bednarek W, Szumiło J, Kotarski J. Accumulation of blood-circulating PD-L1-expressing M-MDSCs and monocytes/macrophages in pretreatment ovarian cancer patients is associated with soluble PD-L1. J Transl Med. 2020;18:220. https://doi.org/10.1186/s12967-020-02389-7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.