

Research

Open Access

Forecasting the cytokine storm following systemic interleukin (IL)-2 administration

Monica C Panelli¹, Richard White², Mareva Foster², Brian Martin³,
Ena Wang¹, Kina Smith¹ and Francesco M Marincola*¹

Address: ¹Immunogenetics Section, Department of Transfusion Medicine, Clinical Center, National Institutes of Health, Bethesda, MD USA, ²Carolinas Medical Center Charlotte, NC USA and ³Unit of Molecular Structure National Institute of Mental Health, National Institutes of Health, Bethesda, MD, USA

Email: Monica C Panelli - mpanelli@mail.cc.nih.gov; Richard White - Richard.White@carolinashealthcare.org; Mareva Foster - Mareva.Foster@carolinashealthcare.org; Brian Martin - martinb@irp.nimh.nih.gov; Ena Wang - Ewang@mail.cc.nih.gov; Kina Smith - Ksmith2@mail.cc.nih.gov; Francesco M Marincola* - fmalincola@mail.cc.nih.gov

* Corresponding author

Published: 02 June 2004

Received: 04 March 2004

Journal of Translational Medicine 2004, **2**:17

Accepted: 02 June 2004

This article is available from: <http://www.translational-medicine.com/content/2/1/17>

© 2004 Panelli et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Extensive clinical experience has shown that systemic interleukin (IL)-2 administration can induce complete or partial regression of renal cell cancer (RCC) metastases in 15 to 20 % of patients. Since IL-2 has no direct anti-cancer effects, it is believed that cancer regression is mediated either by a direct modulation of immune cell effector functions or through the mediation of soluble factors released as a result of IL-2 administration.

We previously observed that transcriptional and protein changes induced by systemic IL-2 administration affect predominantly mononuclear phagocytes with little effect, particularly within the tumor microenvironment, on T cell activation, localization and proliferation. It further appeared that mononuclear phagocyte activation could be best explained by the indirect mediation of a secondary release of cytokines by IL-2 responsive cells either in the circulation or in peripheral tissues.

To better characterize the cytokine outburst that follows systemic IL-2 administration we followed the serum levels of 68 soluble factors in ten patients with RCC undergoing high dose (720,000 IU/kg intravenously every 8 hours) IL-2 therapy. Serum was collected before therapy, 3 hours after the 1st and 4th dose and assayed on a multiplexed protein array platform. This study demonstrated that 1) the serum concentration of more than half the soluble factors studied changed significantly during therapy; 2) changes became more dramatic with increasing doses; 3) subclasses of soluble factors displayed different kinetics and 4) cytokine patterns varied quantitatively among patients.

This study shows that the cytokine storm that follows systemic IL-2 administration is complex and far-reaching inclusive of soluble factors with disparate, partly redundant and partly contrasting effects on immune function. Therefore comparing in parallel large number of soluble factors, it sets a comprehensive foundation for further elucidation of "cytokine storm" in larger patient pools. Based on this analysis, we propose a prospective collection of serum samples in a larger cohort of patients undergoing IL-2 administration with the purpose of discerning patterns predictive of clinical outcome and toxicity.

Background

Interleukin-2 (IL-2) has been extensively used as a single agent for anti-cancer immunotherapy [1,2]. This immune modulatory cytokine can induce long term cures and has been approved by the Food and Drug Administration for standard therapy of metastatic melanoma and renal cell cancer. In addition, it appears that systemic IL-2 administration may increase the frequency of tumor regressions mediated by tumor antigen-specific immunization [3]

Although, it has been postulated that IL-2 acts through the *in vivo* expansion and activation of cytotoxic T lymphocytes [1] and/or the promotion of their migration within target tissues [3], it is likely that at the pharmacologic doses used therapeutically IL-2 has broader immune/pro-inflammatory effects [4,5]. Thus, the mechanism(s) through which systemic IL-2 administration mediates cancer regression remains unknown. It is also unknown whether the substantial toxicity associated with IL-2 administration (that limits its therapeutic usefulness) [6] is mediated through pathways common or distinct from those mediating its anti-cancer effects [7]. It has been observed that hematological toxicity induced by IL-2 (thrombocytopenia and lymphopenia) is associated with increased frequency of clinical responses [8]. In addition, reversal of IL-2-induced toxicity by corticosteroid or anti-TNF- α antibodies blunts its therapeutic effects suggesting a proinflammatory nature for this association [9,10]. Yet, these and other associations between therapeutic and toxic effects are weak and inconclusive and in all probability different pathways may mediate the two phenomena. Obviously, the ability to segregate the anti-cancer effects of IL-2 in target tissues from its systemic toxic effects may be central for the development of improved anti-cancer therapies [11].

We have recently analyzed changes induced by the systemic administration of IL-2 on the transcriptional profile of circulating mononuclear cells and melanoma metastases [12]. This was done by comparing samples obtained before therapy and three hours after IL-2 administration, a time point found to be most informative based on preliminary time course studies. Samples were serially obtained through blood draws and fine needle aspirates of melanoma metastases following a previously validated RNA amplification procedure [13]. This study suggested that the systemic effects of IL-2 administration diverged from those identifiable within the tumor microenvironment since in the latter they seemed primarily restricted to the activation of mononuclear phagocytes. In fact, systemic IL-2 administration had very little direct effects on lymphocyte activation or migration to the tumor site, nor had direct effects on the survival of tumor cells. More specifically, it appeared that IL-2 induced activation of resident mononuclear phagocytes into mature antigen

presenting cells. In addition IL-2 induced activation of cytolytic mechanisms and a massive production of chemo-attractants that could secondarily recruit other immune cells. In addition, the hypothesis that the direct or indirect effects of IL-2 in the target organ are mediated through mononuclear phagocyte activation is in line with the recently reported observation that macrophage activation by IL-2 is essential for tumor regression [14].

In vitro testing failed to identify IL-2 as the direct agonist responsible for the activation of mononuclear macrophages (unpublished observation) suggesting that a down-stream cascade of immune modulators produced by IL-2-responsive cells *in vivo* might be the principal mediator of the effect of this cytokine on the tumor microenvironment. Since it is unlikely that IL-2 directly mediates all of the effects observed after its systemic administration, we hypothesized that the changes in the transcriptional program of melanoma metastases were most likely resulting from an array of soluble factors secondarily produced by IL-2 receptor-bearing immune cells. Indeed, a preliminary *in vitro* analysis of supernatants from cultures of circulating mononuclear cells exposed to IL-2 revealed an intense production of a broad array of cytokines and other soluble factors [15].

The cytokine cascade or cytokine storm induced by systemic IL-2 administration has been extensively reported in the past and attempts have been made to associate individual parameters with biological effects [16,-19]. However, these studies were limited by the number of cytokines tested whose selection was based on an educated guess of their relevance to immune modulation. Recent advances in high-throughput analysis of gene and protein expression allow, however, a global and unbiased overview of complex biological processes adding novel dimensions to the study of this problem. Here we studied the serum levels of 68 soluble factors including cytokines, chemokines, growth factors, soluble receptors and metalloproteinases in patients with RCC undergoing systemic IL-2 therapy. The 68 soluble factors were assembled in a protein array platform (Searchlight™, Boston, MA) without predetermined bias except for their known or suspected relevance to immune phenomena. This was done with two main purposes: the first was to attain a general estimate of the ratio of biologically active factors induced by systemic IL-2 administration out of a pool of randomly chosen reagents. This parameter yields a rough estimate of the complexity and lack of specificity of systemic IL-2 treatment. The second was to provide a map of cytokine secretion patterns relevant to clinical IL-2 administration that could help forecast its toxic or beneficial effects in future prospective studies.

Table 1: Characteristics of the patients enrolled in the study

Patient #	Sampling: pre, post dose#	Course/ Cycle of Sampling	Response	Biological Response	# of doses
p1	pre, post 1	c1c2	SD 2C	NR	3
p2	pre, post 1, post 4	c3c1	SD 2C	NR	8
p3	pre, post 1, post 5	c3c1	PR 1C CR 3C	BR	5
p4	pre, post 1, post 4	c2c1	PR 1C SD 2C PD	BR	7
p5	pre, post 1, post 4	c1c2	SD 1C PD 2C	NR	6
p6	pre, post 1	c1c2	PR 2C PD 3C	BR	4
p7	pre, post 1	c2c1	SD 1C	NR	4
p8	pre	c1c1	PD 1C	NR	8
p9	pre, post 1, post 4	c1c1	PD 1C	NR	7
p10	pre, post 1	c1c2	MR 1C	BR	7
p11	pre, post 1	c3c2	SD 1C PR 3C	NR	3
p12	pre, post 1, post 4	c1c1	SD	NR	8
p13	pre, post 1, post 4	c1c1	PD 1C	NR	7
p14	pre, post 1, post 4	c2c1	MR 1C	BR	8
p15	pre, post 1, post 4	c1c1	MR 1C	BR	7
p16	pre, post 1, post 4	c1c1	PD	NR	9

C = course; CR = complete response; SD = stable disease; PR = partial response; MR = mixed response; PD = progressive disease; BR Biological response = CR+PR+MR; NR = Non responder = PD+SD

Table 2: Toxicity evaluation in the 16 patients with RCC enrolled in this study

Patient #	Age yr	Doses	Hemoglobin	WBC count	Platelet Count	Bilirubin	SGOT	Creatinine
p1	41	3	grade 2	*	*	*	*	*
p2	56	8	*	*	grade 1	*	*	*
p3	53	5	grade 1	*	*	*	*	*
p4	54	7	*	*	grade 1	grade 1	grade 1	grade 1
p5	58	6	grade 2	grade 2	grade 2	grade 3	grade 1	grade 3
p6	47	4	grade 1	*	*	*	*	*
p7	54	4	*	*	*	*	*	grade 2
p8	41	8	grade 1	*	*	grade 2	grade 1	*
p9	44	7	*	*	*	*	grade 1	*
p10	36	7	*	*	*	*	*	*
p11	60	3	*	*	*	*	*	*
p12	45	8	*	*	grade 1	*	grade 1	grade 2
p13	70	7	*	*	*	grade 1	grade 1	*
p14	36	8	*	grade 1	grade 3	*	*	*
p15	66	7	*	*	*	*	grade 3	*
p16	56	9	*	*	*	*	grade 2	*

Toxicity grade is based on the CMC common toxicity criteria

The results suggested that 1) the serum concentration of more than half the soluble factors studied changed significantly during therapy; 2) changes became more dramatic with increasing doses; 3) subclasses of soluble factors displayed different kinetics and 4) cytokine patterns varied quantitatively among patients.

Results and Discussion

Serum was collected from 16 patients (Table 1 and Table 2). From 15 patients, samples could be obtained before treatment and 3 hours after the 1st dose. However, in only 10 patients could the pre-determined collection be completed by obtaining serum before therapy and 3 hours after the 1st and 4th dose of IL-2. Time points of serum collection were selected based on previous experience suggesting incremental effects of IL-2 administration with

Table 3: Soluble factor (pg/ml) concentrations in the serum of 15 patients enrolled in the study (only factors that demonstrated significant changes in concentration after the 1st dose compared with before therapy are shown)

Soluble Factor	Average before IL-2	STD error before	Average after 1 dose	STD error after 1 dose	p value before vs. after 1 dose	Bonferroni correction
IL2*	0	0	3269	841	0.00126	0.07409
TNFR2	3197	222	6215	462	0.00000	0.00001
MMP9	246408	29501	764715	53413	0.00000	0.00008
TNFR1	1659	92	5217	568	0.00000	0.00024
IFN γ	152	17	235	19	0.00000	0.00027
IP10	152	13	2006	267	0.00001	0.00030
MIP3b	151	12	421	45	0.00001	0.00040
MCP2	36	6	1776	261	0.00001	0.00045
MIPIb	525	137	35905	5761	0.00002	0.00100
MMP8	1502	321	3977	537	0.00002	0.00109
SDF1b	75	16	836	137	0.00005	0.00322
Lymph	142	20	338	43	0.00007	0.00407
IL7	3	1	10	2	0.00008	0.00477
IL5	1	1	9	2	0.00012	0.00691
MCP3	12	1	23	3	0.00020	0.01172
I309	22	4	73	12	0.00026	0.01547
E-Selectin	111526	5901	127683	5461	0.00030	0.01753
IL15	1	1	6	1	0.00038	0.02261
Eotaxin	130	16	401	65	0.00047	0.02757
ITAC	31	3	139	25	0.00056	0.03314
MIPIa	3000	111	3228	104	0.00102	0.06033
Gro-A	297	95	1848	388	0.00104	0.06121
Angio2	467	58	671	71	0.00107	0.06330
IL6	61	13	1092	265	0.00112	0.06589
VegF	1112	139	1498	132	0.00130	0.07661
MCP1	1327	248	124449	32058	0.00136	0.08025
MIG	262	35	933	178	0.00262	0.15480
IL8	65	4	800	211	0.00278	0.16379
IL4	0	0	4	1	0.00422	0.24896
TNFA	539	48	640	50	0.00719	0.42432
TARC	282	47	520	88	0.00939	0.55398
MIP3a	17	3	827	291	0.01195	0.70484
IL6R	24605	1759	27935	2652	0.01393	0.82204
IL10	2	1	41	15	0.01517	0.89505
MDC	605	77	661	74	0.03338	1.96915
MMP10	1440	148	1645	147	0.04876	2.87674
MMP1	13937	1775	16437	1602	0.05007	2.95398

Soluble Factors (pg/ml) present in the serum of patients undergoing high dose IL-2 immunotherapy were averaged before and after 1 dose of IL-2. A paired 2-tailed student t test was used to assess significance with an arbitrary cutoff of $p=0.05$. Bonferroni's correction was applied to the data set by correcting for a total of 59 soluble factors (IL-2 was not factored in); IL-2 levels were included as internal control (*) since these values reflect in part administered human recombinant IL-2.

increasing doses [12]. It was felt the four doses represented a reasonable midpoint of IL-2 therapy that could be achieved in most patients before interruption of treatment due to limiting toxicity. Serum collections were obtained 3 hours after IL-2 administration because it is around this time that most of the systemic signs of IL-2 administration are apparent suggesting that the highest levels of soluble factors are reached around that time point. Serum was immediately separated to avoid protein degradation and stored frozen. All serum samples were thawed simultaneously at the completion of the study and immediately tested for the presence of 68 soluble factors (including chemokines, cytokines, soluble cytokine receptors and cell surface molecules) using Searchlight™ arrays.

Cytokines whose concentration in the present conditions was consistently below the threshold of detection of the assay (LIF, IFN α , GM-CSF, IL-1 α , IL-1 β , IL-9, IL-12p70, IL-13) were excluded from the analysis because we could not rule out that significant changes may have occurred with treatment that could not be detected due to the limit of the assay sensitivity. In addition, we excluded IL-2 because it was impossible to distinguish the endogenous levels produced by IL-2 stimulated cells from the circulating exogenously administered recombinant IL-2. The serum concentration of 61 % of evaluable factors changed significantly after one dose of IL-2 (36/59) (Table 3). Similarly, 56 % of the evaluable soluble factors (33/59) increased significantly in concentration after 1 dose of IL-2 in the 10 patients who completed the study (Table 4). After four doses, the concentration of the majority of soluble factors

Table 4: Soluble factor (pg/ml) concentrations in the serum of 10 patients who completed the study (only factors that demonstrated significant changes in concentration after the 1st dose compared with before therapy are shown)

Soluble Factor	Average before IL-2	STD error before IL-2	Average after 1 dose	STD error after 1 dose	p value before vs. after 1 dose	Bonferroni Correction
IL2*	0	0	3668	1178	0.00948	0.55953
TNFR1	1641	112	4702	512	0.00005	0.00292
MMP8	1187	455	3340	500	0.00008	0.00486
TNFR2	3195	259	6014	628	0.00010	0.00588
MCPI	984	227	81270	13486	0.00013	0.00787
IP10	146	18	1571	273	0.00047	0.02778
MCP2	32	9	1500	291	0.00050	0.02950
MMP9	233004	43617	736428	76301	0.00051	0.02984
I309	18	4	52	8	0.00057	0.03372
IFNγ	149	21	222	25	0.00059	0.03477
IL5	0	0	8	2	0.00094	0.05563
MIP1b	512	205	30149	6560	0.00097	0.05714
MIP3b	142	17	347	51	0.00108	0.06345
VegF	865	131	1392	181	0.00175	0.10307
IL7	2	1	10	2	0.00228	0.13444
IL15	0	0	6	2	0.00302	0.17834
E-Selectin	106029	5703	118981	5972	0.00351	0.20688
SDF1b	62	21	646	154	0.00372	0.21974
Lymph	111	22	259	39	0.00400	0.23580
IL6	50	5	747	201	0.00521	0.30754
MCP3	10	1	17	2	0.00616	0.36333
MMPI	12912	2478	17411	2235	0.00686	0.40490
Eotaxin	115	22	318	65	0.00762	0.44948
MIP1a	2952	143	3215	141	0.00790	0.46638
Angio2	494	78	655	101	0.01068	0.63039
MCP4	84	19	264	71	0.01082	0.63834
ITAC	29	4	110	26	0.01104	0.65159
TARC	248	59	478	112	0.02152	1.26942
IL8	61	4	456	151	0.02259	1.33290
IL4	0	0	5	2	0.02601	1.53464
TNFa	521	65	610	59	0.02824	1.66587
Gro-A	314	146	1401	468	0.03637	2.14567
TIMP2	75277	5629	85674	4973	0.04508	2.65947
KGF	373	117	470	97	0.04513	2.66289

Soluble Factors (pg/ml) present in the serum of patients undergoing high dose IL-2 immunotherapy were averaged before and after 1 dose of IL-2. A paired 2-tailed student *t* test was used to assess significance with an arbitrary cutoff of $p=0.05$. Bonferroni's correction was applied to the data set by correcting for a total of 59 soluble factors (IL-2 was not factored in); IL-2 levels were included as internal control (*) since these values reflect in part administered human recombinant IL-2.

studied (76%; 45/59) increased significantly (Table 5). Interestingly, the serum level of 59 % (35/59) of the evaluable soluble factors changes significantly between the 1st and after 4th dose (Table 6). While the serum levels of most soluble factors increased progressively during the 24 hours that spanned between the first and the fourth dose a smaller group including MCP-2, MIP-1 β , MCP-1, NAP-2, IL-6R and Rantes decreased in concentration between the first and the fourth dose. Two factors, MMP-3 and Exod-2 decreased 3 hours after the first dose but increased noticeably after the 4th. Although the results presented in this study are exploratory and fewer cytokine changes maintained significance after correcting the p_2 -value for

the number of tests according to Bonferroni's method [20], it is likely that the changes observed are representative of true biological behavior of these factors in response to systemic IL-2 administration and a more powerful study will confirm these results by allowing a higher level of significance.

We then classified the various factors according to their pattern of expression in response to IL-2 by applying unsupervised hierarchical clustering [21]. We limited this analysis to the 47 cytokines whose concentration levels were significantly altered after the 4th dose (Figure 1). Hierarchical clustering ranks experiments according to

Table 5: Soluble factor (pg/ml) concentrations in the serum of 10 patients who completed the study (only factors that demonstrated significant changes in concentration after the 4th dose compared with before therapy are shown)

Soluble Factor	Average before IL-2	STD error before IL-2	Average after 4 doses	STD error after 4 doses	p value before vs. after 4 doses	Bonferroni correction
IL2*	0	0	1426	394	0.00410	0.24205
MCP2	32	9	296	34	0.00001	0.00047
I309	18	4	222	25	0.00001	0.00070
IP10	146	18	2940	390	0.00003	0.00178
VegF	865	131	1411	132	0.00003	0.00191
ITAC	29	4	438	57	0.00003	0.00200
TNFR1	1641	112	6067	651	0.00004	0.00207
IFN γ	149	21	246	16	0.00004	0.00234
IL15	0	0	14	2	0.00005	0.00269
MCP1	984	227	31267	4583	0.00006	0.00367
MIP3b	142	17	33836	5140	0.00007	0.00405
Lymph	111	22	639	95	0.00010	0.00571
E-Selectin	106029	5703	377934	40503	0.00011	0.00638
TARC	248	59	8993	1517	0.00015	0.00867
SDF1b	62	21	883	142	0.00018	0.01045
MMP8	1187	455	5081	927	0.00024	0.01406
MCP3	10	1	53	8	0.00029	0.01685
MDC	586	111	1272	214	0.00031	0.01841
Eotaxin	115	22	392	67	0.00032	0.01890
TIMP2	75277	5629	108359	4266	0.00033	0.01949
Exodus2	419	71	1108	157	0.00049	0.02911
MMP13	839	221	1742	171	0.00055	0.03224
TIMP1	198383	20136	1021646	178036	0.00071	0.04194
TNFa	521	65	839	74	0.00086	0.05090
MIPIb	512	205	4072	794	0.00105	0.06189
TNFR2	3195	259	16353	3113	0.00115	0.06774
HGF	8281	2359	17298	1311	0.00251	0.14822
IL5	0	0	40	10	0.00274	0.16167
MMP9	233004	43617	657077	142793	0.00391	0.23066
IL7	2	1	6	2	0.00508	0.29968
MMP2	210099	37913	399516	50101	0.00510	0.30111
MMP10	1467	220	3247	528	0.00545	0.32168
MCP4	84	19	1663	495	0.00698	0.41181
IL10	3	1	179	54	0.00707	0.41723
MIP3a	18	4	1346	414	0.00784	0.46274
IL6	50	5	608	184	0.01139	0.67211
VCAM1	310701	30008	1278753	336280	0.01248	0.73618
MMP3	27613	6960	59390	16227	0.01433	0.84561
KGF	373	117	505	109	0.01473	0.86914
ICAM1	499363	62669	2313052	682964	0.01505	0.88787
IL6R	23556	1882	19437	1316	0.02143	1.26450
IL4	0	0	5	2	0.02359	1.39181
IL8	61	4	296	90	0.02410	1.42202
IL18	67	11	167	43	0.03041	1.79442
IL16	13	6	104	43	0.04344	2.56278
MIPIa	2952	143	3160	156	0.05446	3.21343

Soluble Factors (pg/ml) present in the serum of patients undergoing high dose IL-2 immunotherapy were averaged before and after 4 doses of IL-2. A paired 2-tailed student *t* test was used to assess significance with an arbitrary cutoff of $p = 0.05$. Bonferroni's correction was applied to the data set by correcting for a total of 59 soluble factors (IL-2 was not factored in); IL-2 levels were included as internal control (*) since these values reflect in part administered human recombinant IL-2.

their proximity to each other taking into account the entire data set. This unsupervised analysis separated individual samples into three groups that matched the time of collection. In fact, samples obtained before treatment, those obtained after 1 dose and those after 4 grouped together with the exception of the sample obtained from patient 16 after the first dose that clustered with the pre-treatment samples. Analysis of the three categories of sam-

ples suggested that soluble factors in serum of patients treated with IL-2 belong to four categories according to their pattern of expression. A first category demonstrated a slight increase after the first dose and stabilized for the duration of the study (Black bar, Figure 1). A second group was characterized by an increase that occurred only after the fourth dose (Blue bar, Figure 1). A third group increased rapidly after one dose but concentrations

Table 6: Soluble factor (pg/ml) concentrations in the serum of 10 patients who completed the study (only factors that demonstrated significant changes in concentration between the 1st and 4th dose are shown)

Soluble Factor	Average before IL-2	STD error before IL-2	Average after 1 dose	STD error after 1 dose	Average after 4 doses	STD error after 4 doses	pvalue after 1 vs after 4 doses	Bonferroni correction
I309	18	4	52	8	222	25	0.00004	0.00232
MIP3b	18	4	347	51	33836	5140	0.00007	0.00414
TARC	248	59	478	112	8993	1517	0.00015	0.00882
E-Selectin	106029	5703	118981	5972	377934	40503	0.00018	0.01034
Exodus2	419	71	307	26	1108	157	0.00028	0.01628
MDC	586	111	644	109	1272	214	0.00029	0.01740
ITAC	29	4	110	26	438	57	0.00033	0.01935
Lymph	111	22	259	39	639	95	0.00083	0.04905
MCP3	10	1	17	2	53	8	0.00101	0.05956
TIMP1	198383	20136	242215	25530	1021646	178036	0.00116	0.06856
HGF	8281	2359	10836	1791	17298	1311	0.00148	0.08712
IL15	0	0	6	2	14	2	0.00148	0.08719
MMP13	839	221	1081	251	1742	171	0.00163	0.09604
MCP2	32	9	1500	291	296	34	0.00174	0.10287
MIPIb	512	205	30149	6560	4072	794	0.00180	0.10631
TNFR2	3195	259	6014	628	16353	3113	0.00221	0.13028
TNFA	521	65	610	59	839	74	0.00281	0.16587
MCPI	984	227	81270	13486	31267	4583	0.00447	0.26392
NAP2	9479881	2329514	10970212	2240959	8251596	2122243	0.00482	0.28444
IL5	0	0	8	2	40	10	0.00564	0.33291
MMP8	1187	455	3340	500	5081	927	0.00852	0.50244
MMP2	210099	37913	201099	41748	399516	50101	0.00889	0.52449
MCP4	84	19	264	71	1663	495	0.00932	0.54966
IL6R	23556	1882	26254	3045	19437	1316	0.01011	0.59636
ICAM1	499363	62669	501688	90740	2313052	682964	0.01143	0.67431
MMP3	27613	6960	24603	5418	59390	16227	0.01149	0.67778
MMP10	1467	220	1694	218	3247	528	0.01209	0.71314
TIMP2	75277	5629	85674	4973	108359	4266	0.01334	0.78706
VCAM1	310701	30008	335385	17713	1278753	336280	0.01469	0.86698
IPI0	146	18	1571	273	2940	390	0.01571	0.92712
IL10	3	1	45	23	179	54	0.02652	1.56497
IL18	67	11	93	19	167	43	0.03116	1.83873
TNFR1	1641	112	4702	512	6067	651	0.03126	1.84407
IL16	13	6	19	11	104	43	0.04434	2.61591
Rantes	40861	10853	56208	8799	34797	6495	0.04881	2.87988

Soluble Factors levels were averaged across 10 patients after 1 and 4 doses of IL-2. A paired 2-tailed student t test was used to assess significance with an arbitrary cutoff of p= 0.05. Bonferroni's correction was applied to the data set by correcting for a total of 59 soluble factors.

decreased by the fourth dose (Orange bar, Figure 1). Finally, a last group increased in concentration after the first dose and progressively continued to increase after the 4th (Red bar, Figure 1).

The majority of changes in the circulating soluble factors tested occurred after 4 doses of IL-2 (Blue bar Figure 1 and Figure 2). In addition a large cluster of cytokines progressively increased from before therapy throughout IL-2 therapy (Figure 3). We observed several markers of systemic and vascular inflammation whose exact function *in vivo* remains to be elucidated (Figure 4). The soluble form of adhesion molecules ICAM1, V-CAM, E selectin have been reported to be similarly elevated in the serum of chronic renal allograft rejection patients [22], multiple sclerosis and Lupus ([23,24], coronary artery disease [25,26], endothelial vasodilatation [27], immunotherapy of tumor patients with TNF alpha [28] and immunotherapy

of melanoma patients with IL-12 [29] (Figure 5). Our findings may suggest similarly to these reports not only that circulating soluble adhesion molecules play a major role in the inflammatory reaction induced during IL-2 administration [12] but that they may be associated to the toxic effects of IL-2 (blood vessel inflammation, vascular damage, leakage, hypotension).

Metalloproteinases (MMP) 2, 3, 10, 13, tissue inhibitor of metalloproteinase (TIMP1), macrophage derived chemo-attractant (MDC) and thymus and activation regulated chemokine (TARC), Lymphotoxin, secondary lymphoid tissue chemokine (Exodus2), and interferon inducible T cell alpha chemo-attractant (I-TAC) were also on average increased after 4 doses of IL-2. The presence of high levels of MMP is in line with the inflammatory process previously described by us at the genomic level [12] and general outburst of cytokines following IL-2 stimulation. In




Figure 1
Unsupervised Hierarchical clustering (Kendall's Tau) of serum samples from RCC patients obtained before, after 1 and 4 doses of IL-2 (720,000 IU/kg). Hierarchical clustering [21] was applied to the data set encompassing 46 cytokines significantly expressed (excluding IL-2) between before and after 4 doses of IL-2 across 10 serum samples (P2,3,4,5,9,12,13,14,15,16) obtained before, after 1 and after 4 doses of IL-2 (720,000 IU/kg). Values corresponding to soluble factors concentration in pg/ml were transformed in natural log (LN) values, average corrected across experimental samples and displayed according to the central method for display using a normalization factor as recommended by Ross [33]. Patients' serum samples clustered according to the three time points of IL-2 administration. Expression of soluble factors segregated in 4 distinct kinetic profiling. Black bar = soluble factor minimally changing from before to after 4 doses; blue bar = soluble factor expression enhanced at 4 doses only; orange bar = soluble factor expression enhanced after 1 dose; red bar = soluble factor expression enhanced at 1 and 4 doses.

fact, MMP are involved in matrix destruction and regeneration, mononuclear phagocyte migration and function as regulatory proteins by promoting the activation and/or degradation of cytokines. High serum levels of MMP 2 and 9 have been reported in inflammatory diseases of the central nervous system and in particular in MS patients and play a leading role in the evolution of demyelinating

lesions([23]. We could thus hypothesize that the presence of high levels of circulating MMP following IL-2 therapy (after 1 and 4 doses) may be responsible for destruction of endothelial basement membranes, extravasation of mononuclear cells and more generally accumulation of toxic byproducts (Figure 4). In addition, it




Figure 2
Soluble factors increasing after 4 doses of IL-2 Soluble factors increasing after 4 doses of IL-2 clustered in figure 1 (blue bar), were averaged across 10 patients. Values represent concentrations in (pg/ml) ± SD.

suggests a key involvement of mononuclear phagocytes activation in the IL-2-induced inflammatory process [12] since these cells are major producers of MMPs.

Interestingly, tissue inhibitor of metalloproteinase (TIMP1) was also highly expressed after 4 doses. Although the concomitant presence of MMP2 and TIMP1 is not surprising since TIMP1 is the natural inhibitor of MMP2 and MMP9 (which we find particularly increased after 1 and 4 doses), the concurrent dramatic increase of these two factors may reflect an inhibitory feed back mechanism limiting the magnitude of IL-2 induced inflammation. It is also possible that TIMP1 at such high doses may

increase cell proliferation and metabolic activity as previously reported [30].

Elevated MDC and TARC, Lymphotactin, Exodus2 and ITAC supports the presence of a strong migration signal for monocytes, NK cells and T cells as described by us at the RNA level [12] and of a potent mobilization of innate and acquired mechanisms of immune defense. Among the soluble factors that increased significantly after 1 dose and either remained elevated or increased further in concentration after 4 doses (Figure 1 Red bar), were a group of potent inflammatory chemo-attractants for monocytes and lymphocytes (MCP-1, MIP-3, IP-10, I-




Figure 3
Soluble factors increasing after 1 and 4 doses of IL-2 Soluble factors increasing after 1 and 4 doses of IL-2 clustered in figure 1 (red bar), were averaged across 10 patients. Values represent concentrations in (pg/ml) ± SD.

309, stromal derived factor (SDF1)), eosinophils (eotaxin), the soluble TNF R1, the cytokines IL-6, 7 and 8.

The high concentrations of soluble TNFR1 in patients undergoing IL-2 therapy are very striking. Circulating levels of TNF-α and sTNFR1 have been reported to be significantly increased in rheumatoid arthritis patients with amyloidosis as those without [31] (Figure 4) and after IL-2 therapy [34] (Figure 5). Thus this soluble receptor may be yet another important player in enhancing the overall IL-2 inflammatory process. The persistence/increase of these factors throughout the first 24 hours of IL-2 administration (1–4 doses) may suggest that these early response proteins are the true initiator of many of the IL-

2 immune enhancing effects. On the other hand, sustained release/accumulation may be the results of soluble receptor saturation and the initiation of a cascade of toxic byproducts.

An additional finding was the identification of two subgroups of patients who seemed to produce different amount of cytokines in response to IL-2; a smaller group (including patient 9, 13 and 16) appeared to produce less cytokines compared with other patients. This is a finding similar to what was previously noted in a small *in vitro* analysis of PBMC stimulated with IL-2 [15]. This finding if corroborated by larger patient series may have




Figure 4
Soluble Adhesion molecules in serum as markers of systemic and vascular inflammation: correlation with high dose IL-2 immunotherapy and other inflammatory diseases. Increased levels of soluble adhesion molecules (sICAM, sVCAM, sV-Selectin, sTNFR1) and metalloproteinases (MMP2, MMP9) have been described in a variety of inflammatory diseases and believed to act as markers of endothelial cell (EC) activation. Potential explanation for IL-2 effects and toxicity in relation to the effects noted in the reported inflammatory diseases, are listed on the right of this panel.

important implications in separating patients who respond or do not respond to therapy.

The number of patients tested was too small to correlate patterns of expression of various factors with the toxicity experienced or the clinical outcome of their treatment (data not shown).

In summary, this pilot study identified different patterns of production of cytokines during IL-2 therapy that may be of relevance to clinical outcome and toxicity if studied in larger patient populations. In addition, patients' varia-

bility was noted with some patients appearing to be more prone to produce cytokines in response to IL-2. Immune polymorphism(s) affecting individual responses to immune stimulation could be at the basis of this phenomenon [32]. Therefore, further studies should include not only collection of serum samples for serum analysis but also of DNA for genetic analysis. Based on this analysis, we propose a prospective collection of serum samples in a larger cohort of patients undergoing IL-2 administration with the purpose of discerning patterns predictive of clinical outcome and toxicity.




Figure 5
Soluble Adhesion molecules in serum as markers of systemic and vascular inflammation: correlation with high dose IL-2 immunotherapy and other therapies. The presence of increased expression of soluble adhesion molecules (sICAM, sVCAM, sV-Selectin) detected in our study, have also been reported in the serum of patients undergoing TNF alpha and IL-12 immunotherapy. Furthermore, increased level of the inflammatory cytokines (IL-8, IL-6) and soluble TNF receptor (sTNFRI) confirmed previous findings in patients receiving high dose immunotherapy with IL-2.

Materials and Methods

Patients

Sixteen patients with metastatic RCC were recruited at the Carolinas Medical Center (Charlotte, NC 28203) to receive standard high-dose (720,000 IU/Kg) IL-2 (Proleukin, Chiron, Emeryville, CA). Serum was collected prior to treatment from the 16 patients, three hours after the first dose in 15 patients and 3 hours after the first and the fourth dose in 10 patients (Table 1). Toxicities and laboratory abnormalities are reported according to the CMC common toxicity criteria in Table 2.

Serum collection

Fifty to 60 ml of peripheral blood were collected at the bedside in BD Vacutainer™ plus (plastic) serum tubes (BD cat # 367820) before (Pre) and 3–4 hours after 1 (1D) and 4 (4D) doses of IL-2. The complete collection was successful in 10 individuals. Pre and 1D collections were obtained from an additional 5 patients. Serum was immediately separated to avoid protein degradation, stored upright for 10 minutes and centrifuged at room temperature in a horizontal rotor at 3,500 rpm for 8 minutes. The serum phase was then transferred to a 50 ml conical tube (BD, falcon # 352098) pooling serum from 2 to

3 separator tubes and subsequently aliquoted (1 ml/vial) in cryogenic vials (Nunc cat #363401, St Pleasant Prairie, WI). Serum aliquots were snap frozen in dry ice and stored at -80°C until use.

Protein platforms

Protein serum levels were assessed on protein-based platforms (Pierce SearchLight Proteome Arrays, Boston, MA). These arrays consist of multiplexed assays that measure 16 or more proteins per well in standard 96 well plates. The arrays are produced by spotting a 2 × 2, 3 × 3 or 4 × 4 patterns of different monoclonal antibodies into each of a 96-well plate. Following a typical sandwich ELISA procedure, signal is generated using a chemiluminescent substrate. The light produced at each spot in the array is captured by imaging the entire plate with a commercially available cooled CCD camera. The data are reduced using image analysis software that calculates exact values (pg/ml) based on standard curves. Each sample was tested for the following 68 human proteins: monocyte inhibitory protein (MIP)-1α; MIP-1β; monocyte chemotactic protein (MCP)-1/CCL1; MCP-2/CCL8; MCP-3/CCL7; MCP-4/CCL13; macrophage inflammatory protein (MIP)-3β / CCL19; MIP-3α / CCL20; exodus2/CCL21(similar to MIP3α); thymus and activation regulated chemokine (TARC/CCL17); I-309/CCL1; eotaxin/CCL11; macrophage derived chemo-attractant (MDC/CCL22); interferon inducible protein 10 (IP10/CXCL10); stromal derived factor (SDF1)-beta; lymphotactin/XCL1; leukocyte inhibitory factor (LIF); rantes/CCL5; monokine induced by gamma interferon (MIG/CXCL9); ITAC; TNFα; IFNγ; IFNα; GM-CSF; ENA-78/CXCL5; Gro-A/CXCL1; neutrophil activating protein (NAP)-2/CXCL7; vascular endothelial growth factor (VEGF); angiotensin (Angio)-2; ciliary neurotrophic factor (CNTF); fibroblast growth factor (FGF)-basic; keratinocyte growth factor (KGF); hematopoietic growth factor (HGF); heparin binding epidermal growth factor (HB-EGF); platelet derived growth factor PDGF-BB; thymopoietin (Tpo); tissue inhibitor of metalloproteinases (TIMP)1; TIMP2; matrix-metalloproteinase (MMP)-1; MMP-2; MMP-3; MMP-8; MMP-9; MMP-10; MMP-13; interleukin (IL)-1α; IL-1β; IL-2; IL-4; IL-5; IL-6; IL-7; IL-8; IL-9; IL-10; IL-12p70; IL-13; IL-12p40; IL-15; IL-16; IL-18; tumor necrosis factor receptor (TNFR)-1; TNFR2, interleukin-6 receptor (IL-6R); ICAM1; VCAM1; E-Selectin/CD62E; L-Selectin/ CD62L.

Statistical analysis

Eight of these factors (LIF, IFNα, GM-CSF, IL-1α, IL-1β, IL-9, IL-12p70, IL-13), were found to be below the limit of detection in all conditions tested and were removed from the data set prior to statistical analysis because we could not determine with certainty whether changes in their expression occurred below such a threshold. Paired two tailed *t* test was used to determine the level of significance

in serum concentration changes for the remaining 60 soluble factors at different time points. IL-2 was included in the analysis as an internal control. Differences were considered significant at an arbitrary cut off p_2 -value of ≤ 0.05 . Significance was also assessed after correction for the number of tests performed applying the Bonferroni's correction to 59 soluble factors (IL-2 was not considered among the tested variables since it was not possible to discern the respective contribution to serum concentration of exogenously administered or endogenously secreted IL-2).

Relatedness in cytokine expression patterns at different time points of IL-2 treatment was tested with unsupervised by applying Eisen's hierarchical clustering methods [21] to the data set encompassing the 45 cytokines and the 10 samples (P2,3,4,5,9,12,13,14,15,16, Figure 1). This tool ranks experiments according to their proximity to each other taking into account the entire data set.

Acknowledgments

We thank Christine Burns, Amy Wilson and Linda Lavigne at Perbio for their technical support in processing Searchlight™ arrays.

References

- Margolin KA: **Interleukin-2 in the treatment of renal cancer.** *Semin Oncol* 2000, **27**:194-203.
- Atkins MB, Lotze MT, Dutcher JP, Fisher RI, Weiss G, Margolin K, et al.: **High-dose recombinant interleukin-2 therapy for patients with metastatic melanoma: analysis of 270 patients treated between 1985 and 1993.** *J Clin Oncol* 1999, **17**:2105-2116.
- Rosenberg SA, Yang JC, Schwartztruber DJ, Hwu P, Marincola FM, Topalian SL, Restifo NP, Dudley ME, Schwarz SL, Spiess PJ, Wunderlich JR, Parkhurst MR, Kawakami Y, Seipp CA, Einhorn JH, White DE.: **Immunologic and therapeutic evaluation of a synthetic tumor associated peptide vaccine for the treatment of patients with metastatic melanoma.** *Nat Med* 1998, **4**:321-327.
- Cotran RS, Pober JS, Gimbrone MA Jr, Springer TA, Wiebke EA, Gaspari AA, Rosenberg SA, Lotze MT.: **Endothelial activation during interleukin 2 immunotherapy. A possible mechanism for the vascular leak syndrome.** *J Immunol* 1988, **140**:1883-1888.
- Kasid A, Director EP, Rosenberg SA: **Induction of endogenous cytokine-mRNA in circulating peripheral blood mononuclear cells by IL-2 administration to cancer patients.** *J Immunol* 1989, **143**:736-739.
- Lotze MT, Matory YL, Rayner AA, Ettinghausen SE, Vetto JT, Seipp CA, Rosenberg SA.: **Clinical effects and toxicity of interleukin-2 in patients with cancer.** *Cancer* 1986, **58**:2764-2772.
- White RL Jr, Schwartztruber DJ, Guleria A, MacFarlane MP, White DE, Tucker E, Rosenberg SA: **Cardiopulmonary toxicity of treatment with high-dose interleukin-2 in 199 consecutive patients with metastatic melanoma or renal cell carcinoma.** *Cancer* 1994, **74**:3212-3222.
- Phan GQ, Attia P, Steinberg SM, White DE, Rosenberg SA: **Factors associated with response to high-dose interleukin-2 in patients with metastatic melanoma.** *J Clin Oncol* 2001, **19**:3477-3482.
- Papa MZ, Vetto JT, Ettinghausen SE, Mule JJ, Rosenberg SA: **Effect of corticosteroid on the antitumor activity of lymphokine-activated killer cells and interleukin 2 in mice.** *Cancer Res* 1986, **46**:5618-5623.
- Fraker DL, Langstein HN, Norton JA: **Passive immunization against tumor necrosis factor partially abrogates interleukin 2 toxicity.** *J Exp Med* 1989, **170**:1015-1020.
- Marincola FM: **Interleukin-2.** *Biol Ther Cancer Updates* 1994, **4**(3):1-16.

12. Panelli MC, Wang E, Phan G, Puhlmann M, Miller L, Ohnmacht GA, Klein HG, Marincola FM: **Genetic profiling of peripheral mononuclear cells and melanoma metastases in response to systemic interleukin-2 administration.** *Genome Biol* 2002(7):RESEARCH0035.
13. Wang E, Miller L, Ohnmacht GA, Liu E, Marincola FM: **High fidelity mRNA amplification for gene profiling.** *Nat Biotechnol* 2000, **18**:457-459.
14. Masztalerz A, Van Rooijen N, Den Otter W, Everse LA: **Mechanisms of macrophage cytotoxicity in IL-2 and IL-12 mediated tumour regression.** *Cancer Immunol Immunother* 2003, **52**:235-242.
15. Panelli MC, Martin B, Nagorsen D, Wang E, Smith K, Monsurro V, Marincola FM: **A genomic and proteomic-based hypothesis on the eclectic effects of systemic interleukin-2 administration in the context of melanoma-specific immunization.** *Cells Tissues Organs* 788-T1 2004 in press.
16. Boccoli G, Masciulli R, Ruggeri EM, Carlini P, Giannella G, Montesoro E, Mastroberardino G, Isacchi G, Testa U, Calabresi F: **Adoptive immunotherapy of human cancer: the cytokine cascade and monocyte activation following high-dose interleukin 2 bolus treatment.** *Cancer Res* 1990, **50**:5795-5800.
17. Deehan DJ, Heys SD, Simpson W, Herriot R, Broom J, Eremin O: **Correlation of serum cytokine and acute phase reactant levels with alterations in weight and serum albumin in patients receiving immunotherapy with recombinant IL-2.** *Clin Exp Immunol* 1994, **95**:366-372.
18. List J, Moser RP, Steuer M, Loudon WG, Blacklock JB, Grimm EA: **Cytokine responses to intraventricular injection of interleukin 2 into patients with leptomeningeal carcinomatosis: rapid induction of tumor necrosis factor alpha, interleukin-1 beta, interleukin-6, gamma-interferon and soluble interleukin-2 receptor (Mr 55,000 protein).** *Cancer Res* 1992, **52**:1123-1128.
19. Whittington R, Faulds D: **Interleukin-2. A review of its pharmacological properties and therapeutic use in patients with cancer.** *Drugs* 1993, **46**:446-514.
20. Miller RG: *Simultaneous statistical inference* New York: Springer-Verlag; 1981.
21. Eisen MB, Spellman PT, Brown PO, Botstein D: **Cluster analysis and display of genome-wide expression patterns.** *Proc Natl Acad Sci U S A* 1998, **95**:14863-14868.
22. Boratynska M: **[Soluble cell adhesion molecules in chronic renal graft rejection].** *Pol Arch Med Wewn* 1998, **100**:410-418.
23. Correale J, Bassani Molinas ML: **Temporal variations of adhesion molecules and matrix metalloproteinases in the course of MS.** *J Neuroimmunol* 2003, **140**:198-209.
24. Zaccagni H, Fried J, Cornell J, Padilla P, Brey RL: **Soluble adhesion molecule levels, neuropsychiatric lupus and lupus-related damage.** *Front Biosci* 2004, **9**:1654-1659.
25. Blankenberg S, Rupprecht HJ, Bickel C, Peetz D, Hafner G, Tiret L, Meyer J: **Circulating cell adhesion molecules and death in patients with coronary artery disease.** *Circulation* 2001, **104**:1336-1342.
26. Hajilooi M, Sanati A, Ahmadih A, Ghofraniha A, Massoud A: **Circulating ICAM-1, VCAM-1, E-selectin, P-selectin, and TNFalpha in patients with coronary artery disease.** *Immunol Invest* 2003, **32**:245-257.
27. Holmlund A, Hulthe J, Millgard J, Sarabi M, Kahan T, Lind L: **Soluble intercellular adhesion molecule-1 is related to endothelial vasodilatory function in healthy individuals.** *Atherosclerosis* 2002, **165**:271-276.
28. Lieww-a-Fa M, Schalkwijk C, van Hinsbergh VW: **Distinct accumulation patterns of soluble forms of E-selectin, VCAM-1 and ICAM-1 upon infusion of TNFalpha in tumor patients.** *Thromb Haemost* 2003, **89**:1052-1057.
29. Mortarini R, Borri A, Tragni G, Bersani I, Vegetti C, Bajetta E, Pilotti S, Cerundolo V, Anichini A: **Peripheral burst of tumor-specific cytotoxic T lymphocytes and infiltration of metastatic lesions by memory CD8+ T cells in melanoma patients receiving interleukin 12.** *Cancer Res* 2000, **60**:3559-3568.
30. Porter JF, Shen S, Denhardt DT: **Tissue inhibitor of metalloproteinase-1 stimulates proliferation of human cancer cells by inhibiting a metalloproteinase.** *Br J Cancer* 2004, **90**:463-470.
31. Maury CP, Liljestrom M, Laiho K, Tiitinen S, Kaarela K, Hurme M: **Tumor necrosis factor alpha, its soluble receptor I, and -308 gene promoter polymorphism in patients with rheumatoid arthritis with or without amyloidosis: implications for the pathogenesis of nephropathy and anemia of chronic disease in reactive amyloidosis.** *Arthritis Rheum* 2003, **48**:3068-3076.
32. Jin P, Wang E: **Polymorphism in clinical immunology. From HLA typing to immunogenetic profiling.** *J Transl Med* 2003, **1**:8.
33. Ross DT, Scherf U, Eisen MB, Perou CM, Rees C, Spellman P, Iyer V, Jeffrey SS, Van de Rijn M, Waltham M, Pergamenschikov A, Lee JC, Lashkari D, Shalon D, Myers TG, Weinstein JN, Botstein D, Brown PO: **Systematic variation in gene expression patterns in human cancer cell lines.** *Nature Genetics* 2000, **24**:227-235.
34. Locker GJ, Kofler J, Stoiser B, Wilfing A, Wenzel C, Wogerbauer M, Steger GG, Zielinski CC, Mader R, Burgmann H: **Relation of pro- and anti-inflammatory cytokines and the production of nitric oxide in patients receiving high-dose immunotherapy with interleukin-2.** *Eur Cytokine Netw* 2000, **11**(3):391-6.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

