

Commentary

Open Access

Optimizing complement-activating antibody-based cancer immunotherapy: a feasible strategy?

Ester Fonsatti¹, Anna Maria Di Giacomo² and Michele Maio*^{1,2}

Address: ¹Cancer Bioimmunotherapy Unit, Department of Medical Oncology, Centro di Riferimento Oncologico, I.R.C.C.S., 33081 Aviano, Italy and ²Division of Medical Oncology and Immunotherapy, Department of Oncology, University Hospital of Siena, 53100 Siena, Italy

Email: Ester Fonsatti - efonsatti@cro.it; Anna Maria Di Giacomo - immunoterapia@virgilio.it; Michele Maio* - mmaio@cro.it

* Corresponding author

Published: 25 June 2004

Received: 25 May 2004

Journal of Translational Medicine 2004, **2**:21 doi:10.1186/1479-5876-2-21

Accepted: 25 June 2004

This article is available from: <http://www.translational-medicine.com/content/2/1/21>

© 2004 Fonsatti et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Passive immunotherapy with monoclonal antibodies (mAb) targeted to specific tumor-associated antigens is amongst the most rapidly expanding approaches to biological therapy of cancer. However, until now a limited number of therapeutic mAb has demonstrated clinical efficacy in selected neoplasia. Results emerging from basic research point to a deeper characterization of specific biological features of neoplastic cells as crucial to optimize the clinical potential of therapeutic mAb, and to identify cancer patients who represent the best candidates to antibody-based immunotherapy. Focus on the tissue distribution and on the functional role of membrane complement-regulatory proteins such as Protectin (CD59), which under physiologic conditions protects tissues from Complement (C)-damage, might help to optimize the efficacy of immunotherapeutic strategies based on C-activating mAb.

Introduction

In view of their potentiality to specifically target tumor cells, antibody-based therapeutic strategies still represent very attractive clinical approaches to cancer treatment. A renewed interest in this field of cancer immunotherapy has derived from the significant clinical results obtained with anti-CD20 and anti-HER2 monoclonal antibodies (mAb) in B cell malignancies and breast cancer, respectively, and from the availability of new molecular techniques such as recombinant DNA technology. Currently, several mAb targeted to different tumor-associated antigens (TAA) are employed in initial clinical studies of passive immunotherapy for solid and hematological malignancies [1].

The tumoricidal activity of therapeutic antibodies depends on different mechanisms of action such as inhibition of downstream signaling events in the target cells.

In addition, therapeutic mAb can directly induce cell death by triggering apoptosis or through "indirect" immunologic mechanisms such as antibody-dependent cell-mediated cytotoxicity (ADCC) and/or Complement (C)-mediated cytotoxicity (CDC).

Over the past years, C-activating mAb have been extensively utilized for the treatment of patients with tumors of different histotype, in particular in patients with cutaneous melanoma which represents a "model disease" to design and activate approaches of biological therapy in cancer patients. Among therapeutic mAb that mediate CDC and ADCC of target melanoma cells, anti-GD2 and anti-GD3 mAb have been largely used in the clinical setting, although with rather unsatisfactory results [2-4]. Nevertheless, in light of the promising results obtained in pre-clinical studies, selected engineered C-activating mAb are still utilized in clinical trials. Along this line, results of

a phase I pilot clinical trial of human IgM mAb directed to GM3 ganglioside in 9 patients with metastatic melanoma have been most recently reported [5].

Several *in vitro* and *in vivo* studies, focused on the mechanisms regulating C-activity and tumor-host interactions, have provided insights on distinct biological features of neoplastic cells that may affect the clinical efficacy of passive immunotherapy with antibodies or their derived molecules [6]. Among the different mechanisms of immune escape triggered by neoplastic cells, the expression of high levels of the C-regulatory proteins such as Protectin (CD59) is emerging as an important strategy that limits the potential clinical benefits deriving from antibody-based immunotherapeutic approaches.

Discussion

The long-standing field of cancer immunotherapy clearly needs well-substantiated pre-clinical evidences on the fine modality of action of the different strategies utilized for cancer treatment. In fact, a variety of therapeutic tools, including C-activating mAb have been extensively utilized in the clinic [2-4], in the absence of accompanying studies designed to fully explore their biologic, functional and clinical potential. As a direct consequence of this partially "blind" therapeutic approach, several potentially useful immunotherapeutic agents and approaches have been rapidly dismissed, due to their limited clinical efficacy. Indeed, opposite to chemotherapy, a much broader number of variables must be taken into account to maximize the clinical accomplishment of immunobiologic agents used as therapeutic tools in cancer patients.

As far as the clinical efficacy of passive immunotherapy of cancer patients with therapeutic mAb that mediate the activation of the C cascade, a major limitation is certainly represented by the presence of a functional form of the C-regulatory protein CD59 on the cell membrane of neoplastic cells, as well as in the tumor microenvironment [7,8]. Among solid tumors, the expression and functional role of CD59 has been well investigated in melanoma. CD59 is broadly expressed on normal and neoplastic tissues of melanocytic origin, with few non-CD59 expressing melanomas, has a limited intra- and inter-lesional heterogeneity and, among known C-regulatory proteins, it represents the main restriction factor of C-susceptibility of human melanomas [9,10]. Furthermore, a statistically significant ($r = 0.914$; $p < 0.001$) inverse correlation was identified between absolute levels of CD59 molecules expressed on melanoma cells and their susceptibility to C-mediated cytotoxicity induced by anti-GD3 mAb R24 [11]. Thus, melanomas from distinct patients were highly susceptible to C-mediated cytotoxicity, while neoplastic cells from other individuals were completely resistant to C-cytotoxicity, even in the presence of elevated amounts

of C-sensitizing mAb R24 [11]. Conversely, no significant correlation was found between levels of cell membrane GD3 expressed on melanoma cells and their C-susceptibility [11]. Thus, in spite of the efficient and rapid tumor targeting potential of therapeutic mAb, these findings strongly caution that their therapeutic efficacy may be greatly implemented through the treatment of patients bearing melanoma lesions that express weak to intermediate levels of CD59 and, thus, are more susceptible to the cytolytic effect of autologous C activated by therapeutic mAb. This notion is clearly not restricted to human melanomas but rather represents a more general phenomenon; in fact, resistance to rituximab (anti-CD20)-mediated C cytotoxicity is associated with high levels of CD59 expression on neoplastic cells of non-Hodgkin's lymphoma and multiple myeloma patients [12].

A soluble form of CD59 (sCD59) has been identified in body fluids and in the culture supernatants of normal and neoplastic cells [8,13,14]. Concerning melanoma, it was demonstrated that the amounts of sCD59 released from CD59-positive melanoma cells correlate with the levels of CD59 expressed on the cell surface [8]. Noteworthy, sCD59 released from melanoma cells blocks the binding of anti-CD59 mAb to cell surface CD59 neutralizing their functional activity [8]. In light of these evidences, the recent observation that CD59 acts as a triggering co-receptor for Natural Killer (NK) cell-mediated cytotoxicity [15] identifies an additional protective role for CD59 in natural immunity. In fact, high amounts of circulating sCD59 detectable in sera of cancer patients may bind to NK cells affecting CD59-mediated functional activation.

Conclusions

Much remains to be gained to make full use of the potential of immunotherapeutic agents in the clinical setting; nevertheless, we are ever more generating pre-clinical and clinical data that allow us to design prospective trials taking into strict account the characterization of the "biological eligibility" of cancer patients to specific therapeutic options, through the analysis of defined phenotypic characteristics of their malignant lesions. Such a strategy will ultimately lead to the identification and selection of patients who represent the best candidates to specific immunotherapeutic approaches.

Abbreviations

Antibody-dependent cell-mediated cytotoxicity, ADCC

Complement, C

Complement-mediated cytotoxicity, CDC

Monoclonal antibodies, mAb

Natural Killer, NK

References

- Harris M: **Monoclonal antibodies as therapeutic agents for cancer.** *Lancet Oncol* 2004, **5**:292-302.
- Irie RF, Morton DL: **Regression of cutaneous metastatic melanoma by intralesional injection with human monoclonal antibody to ganglioside GD2.** *Proc Natl Acad Sci USA* 1986, **83**:8694-8698.
- Soiffer RJ, Chapman PB, Murray C, Williams L, Unger P, Collins H, Houghton AN, Ritz J: **Administration of R24 monoclonal antibody and low-dose interleukin 2 for malignant melanoma.** *Clin Cancer Res* 1997, **3**:17-24.
- Scott AM, Lee FT, Hopkins W, Cebon JS, Wheatley JM, Liu Z, Smyth FE, Murone C, Sturrock S, MacGregor D, Hanai N, Inoue K, Yamasaki M, Brechbiel MW, Davis ID, Murphy R, Hannah A, Lim-Joon M, Chan T, Chong G, Ritter G, Hoffman EW, Burgess AW, Old LJ: **Specific targeting, biodistribution, and lack of immunogenicity of chimeric anti-GD3 monoclonal antibody KM871 in patients with metastatic melanoma: results of a phase I trial.** *J Clin Oncol* 2001, **19**:3976-3987.
- Irie RF, Ollila DW, O'Day S, Morton DL: **Phase I pilot clinical trial of human IgM monoclonal antibody to ganglioside GM3 in patients with metastatic melanoma.** *Cancer Immunol Immunother* 2004, **53**:110-117.
- Gelderman KA, Tomlinson S, Ross GD, Gorter A: **Complement function in mAb-mediated cancer immunotherapy.** *Trends Immunol* 2004, **25**:158-164.
- Gelderman KA, Hakulinen J, Hagensaars M, Kuppen PJ, Meri S, Gorter A: **Membrane-bound complement regulatory proteins inhibit complement activation by an immunotherapeutic mAb in a syngeneic rat colorectal cancer model.** *Mol Immunol* 2003, **40**:13-23.
- Brasoveanu LI, Fonsatti E, Visintin A, Pavlovic M, Cattarossi I, Colizzi F, Gasparollo A, Coral S, Horejsi V, Altomonte M, Maio M: **Melanoma cells constitutively release an anchor-positive soluble form of protectin (CD59) that retains functional activities in homologous complement-mediated cytotoxicity.** *J Clin Invest* 1997, **100**:1248-1255.
- Junnikkala S, Hakulinen J, Meri S: **Targeted neutralization of the complement membrane attack complex inhibitor CD59 on the surface of human melanoma cells.** *Eur J Immunol* 1994, **24**:611-615.
- Maio M, Brasoveanu LI, Coral S, Sigalotti L, Lamaj E, Gasparollo A, Visintin A, Altomonte M, Fonsatti E: **Structure, distribution, and functional role of protectin (CD59) in complement-susceptibility and in immunotherapy of human malignancies.** *Int J Oncol* 1998, **13**:305-318.
- Brasoveanu LI, Altomonte M, Fonsatti E, Colizzi F, Coral S, Nicotra MR, Cattarossi I, Cattelan A, Natali PG, Maio M: **Levels of cell membrane CD59 regulate the extent of complement-mediated lysis of human melanoma cells.** *Lab Invest* 1996, **74**:33-42.
- Treon SP, Mitsiades C, Mitsiades N, Young G, Doss D, Schlossman R, Anderson KC: **Tumor cell expression of CD59 is associated with resistance to CD20 serotherapy in patients with B-cell malignancies.** *J Immunother* 2001, **24**:263-271.
- Rooney IA, Atkinson JP, Krul ES, Schonfeld G, Polakoski K, Saffitz JE, Morgan BP: **Physiologic relevance of the membrane attack complex inhibitory protein CD59 in human seminal plasma: CD59 is present on extracellular organelles (protasomes), binds cell membranes, and inhibits complement-mediated lysis.** *J Exp Med* 1993, **177**:1409-1420.
- Bjorge L, Vedeler CA, Ulvestad E, Matre R: **Expression and function of CD59 on colonic adenocarcinoma cells.** *Eur J Immunol* 1994, **24**:1597-1603.
- Marcenaro E, Augugliaro R, Falco M, Castriconi R, Parolini S, Sivori S, Romeo E, Millo R, Moretta L, Bottino C, Moretta A: **CD59 is physically and functionally associated with natural cytotoxicity receptors and activates human NK cell-mediated cytotoxicity.** *Eur J Immunol* 2003, **33**:3367-3376.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

